

## UGC-MRP PROJECT REPORT

### **“CHARACTERIZATION OF JUVENILE HORMONE ANALOGUE-RESPONSIVE PROTEIN FROM THE LARVAL HAEMOLYMPH OF *SPODOPTERA MAURITIA* (LEPIDOPTERA: NOCTUIDAE)”**

Principal investigator: Dr. Kannan V.M.

Professor, Department of Zoology

University of Calicut. [kannanvm@yahoo.com](mailto:kannanvm@yahoo.com)

## INTRODUCTION

Insects are the largest group in the animal kingdom. Some of them are pests and cause considerable economic loss. *Spodoptera mauritia* or rice swarming caterpillar is a sporadic pest of paddy. It is estimated that the loss in yield caused by larval infestation of *Spodoptera mauritia* range from 10 to 20%. In India it is found in all the rice growing areas especially along the west coast and delta in Kerala and Tamil Nadu (David and Ananthakrishnan, 2004). Pest management is an integral part of any successful agriculture. Application of conventional insecticides poses great threat to the human health and environment. Development of more eco-friendly pest management approaches is of prime importance for food security and health. Use of Insect Growth Regulators (IGRs) for pest management is an alternative as they are target-specific, non-persistent, biodegradable and environmentally benign substances, with less toxicity to non-target organisms. Many IGR's are juvenile hormone or ecdysone agonists.

Protein metabolism plays a key role in rebuilding adult structures during the transformation of larvae/pupae into adult. Hemolymph protein levels generally increase during each instar but decline during moulting. Several hemolymph proteins like insect hexamerins are thought to transport hormones, phenols /or some cuticular proteins to the hypodermis. Typically two to four physico-chemically distinct storage protein species occur. The last larval instar of holometabolous insects has been characterized by active synthesis of arylphorin (aromatic amino acids bearing storage proteins) and pupal storage proteins (Wyatt, 1978) During metamorphosis larval plasma proteins were hydrolyzed to free amino acids and major part being incorporated into new adult proteins. Thus haemolymph proteins are crucial for insect development. In this

study we demonstrate the effect of pyriproxyfen, a Juvenile hormone analogue on larval protein profile of *Spodoptera mauritia* Boisd. (Lepidoptera: Noctuidae) or paddy army worm, a pest of paddy (*Oriza sativa*).

### **OBJECTIVES OF THE PROJECT (AS GIVEN IN THE PROPOSAL)**

1. To identify and characterize the JH analogue-responsive proteins.
2. To identify the site of synthesis of JH analogue-responsive proteins.
3. To understand the regulation of the JH analogue-responsive proteins by JH analogues.

### **STANDARDIZATION OF THE CULTURING TECHNIQUES OF *SPODOPTERA MAURITIA***

We standardized the conditions for culturing *Spodoptera mauritia*. The adult moths were collected using fluorescent light traps. They were kept in glass beakers and fed with a dilute solution of honey. They were allowed to mate and lay eggs. The egg will hatch within 3-5 days. The larvae hatched out were fed with fresh leaves of the grass *Ischaemum aristatum* and were maintained at room temperature (28°C.)

### **TOXICITY OF PYRIPROXYFEN TO LARVAE OF *SPODOPTERA MAURITIA***

The average percentage mortality for 3<sup>rd</sup>, 4<sup>th</sup>, 5<sup>th</sup> & 6<sup>th</sup> instar larvae of *Spodoptera mauritia* treated with different concentrations of pyriproxyfen was calculated (Table 1.) With increase in concentration of pyriproxyfen, the mortality increased in all the instars of larvae tested.

Amount of pyriproxyfen applied/ larva	Average percentage mortality $\pm$ SE			
	3 <sup>rd</sup> instar	4 <sup>th</sup> instar	5 <sup>th</sup> instar	6 <sup>th</sup> instar
Control	0	0	0	0
5 $\mu$ g	14.4 $\pm$ 2.1	12.5 $\pm$ 2.5	.....	.....
10 $\mu$ g	25 $\pm$ 5.0	27.5 $\pm$ 6.4	.....	.....
25 $\mu$ g	85 $\pm$ 2.9	70.8 $\pm$ 5.1	15.8 $\pm$ 2.0	.....
50 $\mu$ g	96.7 $\pm$ 3.3	91.7 $\pm$ 4.4	65.8 $\pm$ 2.2	.....
100 $\mu$ g	100 $\pm$ 0.0	100 $\pm$ 0.0	75 $\pm$ 2.9	8.13 $\pm$ 2.8
125 $\mu$ g	.....	.....	96.7 $\pm$ 3.3	13.8 $\pm$ 1.3
200 $\mu$ g	.....	.....	.....	32.5 $\pm$ 2.5
300 $\mu$ g	.....	.....	.....	45 $\pm$ 5.0
400 $\mu$ g	.....	.....	.....	73.3 $\pm$ 1.7

**Table I:** Percentage mortality of 3<sup>rd</sup> 4<sup>th</sup> 5<sup>th</sup> & 6<sup>th</sup> instar larvae of *Spodoptera mauritia* treated with different concentrations of pyriproxyfen

## CALCULATION OF LD<sub>50</sub> VALUE

The LD<sub>50</sub> value (24 hours) of pyriproxyfen for the 3<sup>rd</sup>, 4<sup>th</sup>, 5<sup>th</sup> & 6<sup>th</sup> instar larvae of *Spodoptera mauritia* was found out by treatment with different concentrations of pyriproxyfen in acetone and the values were 14.13 $\pm$ 2.67, 15.85 $\pm$ 3.67, 39.81 $\pm$ 2.61 and 316.20 $\pm$ 2.64  $\mu$ g/larvae respectively (Table 2)

SL. NO.	LARVAL INSTAR	LD <sub>50</sub> VALUE (µg) /larva (MEAN ±SE)
1	THIRD	14.13±2.67
2	FOURTH	15.83±3.67
3	FIFTH	39.81±2.61
4	SIXTH	316.20±2.64

**Table 2:** LD<sub>50</sub> value (24 hours) of pyriproxyfen for 3<sup>rd</sup> 4<sup>th</sup> 5<sup>th</sup> & 6<sup>th</sup> instar larvae of *Spodoptera mauritia*

## EFFECT OF PYRIPROXYFEN ON HAEMOLYMPH PROTEIN CONCENTRATION

In our initial study we found that treatment of 5<sup>th</sup> instar larvae with sub lethal concentrations of pyriproxyfen on day 0, led to a statistically significant ( $p < 0.05$ ) increase in total haemolymph protein concentration estimated by modified Lowry's method (Sandermann and H, Jr, Stromiger 1972) on day 1 after treatment, compared to control

SL. No.	SAMPLE	AMOUNT OF PROTEIN (µg/µl) ± SE
1	Control	3.02±0.02
2	LD <sub>10</sub>	3.23±0.03

**Table 3:** The difference in haemolymph protein concentration on treatment of pyriproxyfen

## EFFECT OF PYRIPROXYFEN ON HAEMOLYMPH PROTEIN PROFILE

Fifth instar larvae were treated with sub lethal concentration ( $LD_{10}$ ) of pyriproxyfen (on day 0) and haemolymph collected after 24 hours of the treatment. Haemolymph was subjected to SDS-PAGE under reducing conditions (Laemmli UK, 1970). The band intensity of an 83 kDa protein band is increased significantly compared to that of control (Fig. 1). This is a juvenile hormone analogue (pyriproxyfen) responsive protein.




Fig. 1: SDS-PAGE (10%) Gel electrophoresis of haemolymph (3 $\mu$ l) of *Spodoptera mauritia* 5<sup>th</sup> Instar larvae.

## DETERMINATION OF GLYCOSYLATION STATUS OF PYRIPROXYFEN-RESPONSIVE PROTEIN




Fig. 2: PAS stained SDS-PAGE (10%) Gel electrophoresis of haemolymph of *Spodoptera mauritia* larvae

The glycosylation status of the identified pyriproxyfen-responsive protein was determined using Periodic Acid-Schiff's (PAS) staining of the SDS-PAGE separated haemolymph proteins (Ref). The band corresponding to the Pyriproxyfen responsive protein (83 kDa) was seen in reddish pink colour which indicates that the identified Pyriproxyfen responsive protein is a glycoprotein.

## IDENTIFICATION OF THE SITE OF SYNTHESIS/STORAGE OF PYRIPROXYFEN-RESPONSIVE PROTEIN

The fat body extract of *Spodoptera mauritia* was subjected to 10% SDS-PAGE along with the haemolymph collected from the control larvae. We found a band in the fat body extract which corresponds to the identified Pyriproxyfen responsive protein in molecular weight.

It is possible that the protein identified in fat body may be the JH-analogue responsive protein found in haemolymph. Thus this protein is expressed /localized in fat body. Hence it is possible that one of the sites of synthesis/storage of the identified Pyriproxyfen- responsive protein may be the fat body.




Fig. 3: SDS-PAGE (10%) Gel electrophoresis of Fat body extract and haemolymph of *Spodoptera mauritia* larvae.

### **DETERMINATION OF THE REGULATION OF THE JH ANALOGUE-RESPONSIVE PROTEIN BY PYRIPROXYFEN**

Fifth instar larvae of *S.mauritia* was treated with different concentrations (10µg, 25µg and 100µg) of pyriproxyfen to determine its role in the regulation of the identified protein. The haemolymph of the treated larvae were collected after 24 hours and subjected to SDS-PAGE. The intensity of the identified protein band was increased with the increasing concentration of pyriproxyfen.




Fig. 4: SDS-PAGE (10%) Gel electrophoresis haemolymph of *Spodoptera mauritia* larvae treated with different concentration of pyriproxyfen

The concentrations of three storage proteins (SL-1,SL-2 and SL-3, hexamers of 70–80 kDa subunits) and two biliverdin-binding proteins (BP-A and BP-B, dimers of 165 kDa) in the haemolymph and fat body during larval and pupal development of *Spodoptera litura* were determined by immunodiffusion tests using polyclonal antisera (ToyoshiYoshiga et al 1997). All these three storage proteins have molecular sizes between 400 and 450 kDa, and are composed of subunit(s) which range in size from 70 to 80 kDa. ( Tojo S and Yoshiga T , 1994). Thus the JH-analogue responsive protein reported from the hemolymph of *Spodoptera mauritia* is likely to be a hexamerine, which is a storage protein in insects. The storage proteins are metabolized and the amino acids are utilized for making the new tissues during development. Some of them also bind and transport ligands including hormones. These proteins are over expressed in response to JH analogue (pyriproxyfen) exposure and it will be worth examining the physiological significance of this insecticide induced over expression.

## CONCLUSIONS

- 1) We determined the average percentage mortality of 3<sup>rd</sup>, 4<sup>th</sup>, 5<sup>th</sup> and 6<sup>th</sup> instar larvae of *Spodoptera mauritia* for different concentrations of Pyriproxyfen (JH-analogue).



- 2) We determined the LD<sub>50</sub> of Pyriproxyfen for 3<sup>rd</sup>, 4<sup>th</sup>, 5<sup>th</sup> and 6<sup>th</sup> instar larvae of *Spodoptera mauritia*.
- 3) Also we found that exposure of 5<sup>th</sup> instar larvae to sub lethal concentration of pyriproxyfen led to an increase in haemolymph protein concentration.
- 4) In addition we showed that the treatment with sub lethal (LD<sub>10</sub>) concentration of pyriproxyfen to 5<sup>th</sup> instar larvae leads to increase in intensity of an 83 kDa protein band.
- 5) The identified JH analogue- responsive protein is a glycoprotein with a subunit molecular mass of 83 kDa.
- 6) Fat body may be one of the sites of synthesis of the Pyriproxyfen-responsive 83 kDa protein.
- 7) The native Pyriproxyfen-responsive protein has subunit with a molecular weight of 83kDa.
- 8) The Pyriproxyfen-responsive protein may be a member of hexamerin family of proteins, which are storage proteins in the haemolymph of insects with subunit molecular weight in the range of 70-80kDa.
- 9) Hexamerins are major proteins in the haemolymph and they have roles in rebuilding adult structures. But these proteins are over expressed in response to JH analogue (Pyriproxyfen) exposure.
- 10) As storage proteins are crucial for insect development, it will be worth examining the physiological significance of this insecticide induced over expression of this protein.

## **PAPERS PUBLISHED**

A part of the work is published in

1. International Journal of Agriculture Innovations and Research (2016) Volume 5, Issue 1, ISSN (Online) 2319-1473.

## **PAPERS PRESENTED IN CONFERENCES**

1. Part of the results obtained in the study is presented in the Fourth Biopesticide international conference (BIOCICON-2013) held at Palayamkottai, Tamil Nadu
2. A part is also presented in the international symposium on Comparative endocrinology and integrative physiology (CEIP-2015) held at Thiruvananthapuram, Kerala

## **AWARDS OBTAINED**

1. 'Best paper award' for the presentation in the Fourth Biopesticide international conference (BIOCICON-2013) held at Palayamkottai, Tamil Nadu.

## **PAPERS COMMUNICATED**

1. The Juvenile Hormone mimic, Pyriproxyfen, increases the level of the major haemolymph protein in the larvae of *Spodoptera mauritia* Bois. (Lepidoptera: Noctuidae) Resmitha C and Kannan Vadakkadath Meethal.

## **REFERENCES**

1. David B. V. And Anantha Krishnan T. N. (2004) General and applied entomology, Tata Mc Graw - Hill Publishing company Limited New Delhi, pp 1184.
2. Laemalli UK, Nature. 1970, 227:680–685.
3. Sandermann, H,Jr, Stromiger JL. J. Biol. Chem, 1972, 247:5123-5131.
4. Toyoshi Yoshiga , Kousei Maruta and Sumio Tojo 1997. J.Insect Physio. 67-76
5. Tojo S, Yoshiga T, Insect Biochemistry and Molecular Biology. 1994, 24: 729-738
6. Wyatt, GR, Pan, ML. A rev. biochem. 1978, 47: 779-817

Dr. Kannan V.M. (PI- UGC MRP)  
Professor, Dept. of Zoology,  
University of Calicut.

**EXECUTIVE SUMMARY OF THE UGC-MRP PROJECT**  
**“CHARACTERIZATION OF JUVENILE HORMONE ANALOGUE-  
RESPONSIVE PROTEIN FROM THE LARVAL HAEMOLYMPH OF  
*SPODOPTERA MAURITIA* (LEPIDOPTERA: NOCTUIDAE)”**

Principal investigator: Dr. Kannan V.M.

Professor, Department of Zoology

University of Calicut. [kannanvm@yahoo.com](mailto:kannanvm@yahoo.com)

1. Determined the average percentage mortality of 3<sup>rd</sup>, 4<sup>th</sup>, 5<sup>th</sup> and 6<sup>th</sup> instar larvae of *Spodoptera mauritia* for different concentrations of Pyriproxyfen (JH-analogue).
2. Determined the LD<sub>50</sub> of Pyriproxyfen for 3<sup>rd</sup>, 4<sup>th</sup>, 5<sup>th</sup> and 6<sup>th</sup> instar larvae of *Spodoptera mauritia*.
3. Also found that exposure of 5<sup>th</sup> instar larvae to sub lethal concentration of pyriproxyfen led to an increase in haemolymph protein concentration.
4. In addition showed that the treatment with sub lethal (LD<sub>10</sub>) concentration of pyriproxyfen to 5<sup>th</sup> instar larvae leads to increase in intensity of an 83 kDa protein band.
5. The identified JH analogue- responsive protein is a glycoprotein with a subunit molecular mass of 83 kDa.
6. Fat body may be one of the sites of synthesis of the Pyriproxyfen-responsive 83 kDa protein.
7. The native Pyriproxyfen-responsive protein has subunit with a molecular weight of 83kDa.
8. The Pyriproxyfen-responsive protein may be a member of hexamerin family of proteins, which are storage proteins in the haemolymph of insects with subunit molecular weight in the range of 70-80kDa.
9. Hexamerins are major proteins in the haemolymph and they have roles in rebuilding adult structures. But these proteins are over expressed in response to JH analogue (Pyriproxyfen) exposure.
10. As storage proteins are crucial for insect development, it will be worth examining the physiological significance of this insecticide induced over expression of this protein.

# Toxicity of Insect Growth Regulator, Pyriproxyfen, on larvae of *Spodoptera mauritia* Bois (Lepidoptera: Noctuidae)

**C. Resmitha**

University of Calicut,

Thenjipalam (PO), Malappuram-673635, Kerala, India

E-mail: [rechujan@gmail.com](mailto:rechujan@gmail.com)

**Vadakkadath Meethal**

University of Calicut,

Thenjipalam (PO), Malappuram-673635, Kerala, India

E-mail: [kannanvm@yahoo.com](mailto:kannanvm@yahoo.com)

\*Author for correspondence

**Abstract** – Insect growth regulators belong to a class of compounds which interfere with normal growth, development and reproduction in insects. Many insect growth regulators are analogues or mimics of insect hormone ecdysone or juvenile hormone. Pyriproxyfen is an insect growth regulator which mimics the action of juvenile hormone. The effects of insect growth regulators on many lepidopteran pests have been studied with respect to its toxicity and developmental alterations. But the toxicity of pyriproxyfen to larvae of *Spodoptera mauritia* is not explored. In this study we treated 3<sup>rd</sup>, 4<sup>th</sup>, 5<sup>th</sup> & 6<sup>th</sup> instar larvae of *Spodoptera mauritia* Bois. with different concentrations of pyriproxyfen (Knack IGR). It was found that the LD<sub>50</sub> value of pyriproxyfen for the 3<sup>rd</sup>, 4<sup>th</sup>, 5<sup>th</sup> and 6<sup>th</sup> instar larvae were 14.13±2.67, 15.85±3.67, 39.81±2.61 and 316.20±2.64 µg/larvae respectively. Studies are ongoing to understand the effect of sub lethal concentrations of pyriproxyfen on the development of *S.mauritia*.

**Keywords** – JH Analogue, Insect Growth Regulators, Pyriproxyfen, *Spodoptera mauritia*, LD<sub>50</sub>.

## I. INTRODUCTION

Insects are the largest group in the animal kingdom. Some of them are pests and cause considerable economic loss. *Spodoptera mauritia* Bois. (Lepidoptera: Noctuidae) or rice swarming caterpillar or paddy army worm is a sporadic pest of paddy distributed all over the world. *Spodoptera mauritia* has six larval instars before pupation. The larvae feed on leaves of paddy or alternate host plant such as *Ischaemum aristatum*. In India it is found in all the rice growing areas especially along the west coast and delta in Kerala and Tamil Nadu. [1]. It has attained the status of a major pest of rice in Eastern India, especially in Orissa, Chhattisgarh, Jharkhand and Bihar.

Pest management is an integral part of any successful agriculture. Application of conventional insecticides poses great threat to the human health and environment. Development of more eco-friendly pest management approaches is of prime importance for human health and environment. Use of Insect Growth Regulators (IGRs) for pest management is an alternative as they are more target-specific, non-persistent, biodegradable and environmentally benign substances, with less toxicity to non-target organisms. Insect Growth Regulators belong to a group of compounds which interfere with normal growth, development and reproduction in insects by disrupting hormonally regulated physiological processes

of insects. Several such compounds are known and their effects on metamorphosis and reproduction in a number of insect species have been extensively studied.[2,3,4] Advantages of Juvenile hormone analogues (JHAs) include their fast penetrance through the insect cuticle, needed only in low concentrations and get degraded to non toxic compounds in a short time period. Already more than 500 analogues with juvenile hormone (JH) activity have been discovered. Among the well known JHAs are, Epofenonane, Methoprene, Hydroprene, Kinoprene, and Fenoxycarb.[5] The earlier reported JHA of commercial success were Methoprene and Hydroprene.[6] Methoprene is active against dipteran insects and fleas and hydroprene is active against cockroach. These compounds however, were too unstable under field conditions to be used for agriculture.[7] Krysan, J. L. (1990) reported the photostable JH analogue, fenoxycarb was effective not only on household pests but also on agricultural pests such as leaf rollers and codling moth.

Pyriproxyfen is a JH analogue preventing the larvae from developing in to adulthood and thus rendering them unable to reproduce. Most common morphogenetic effect of JHA treatment is the production of extra larval, nymphal or pupal form. The formation of extra larval instar depends on stage and age of the larvae at the time of treatment. Pyriproxyfen has relatively low mammalian toxicity and was first registered in Japan in 1991 for controlling public health pests.[8] Pyriproxyfen is a commonly used insect growth regulator against whiteflies.[9,10] In 1996 it was introduced in US to protect cotton crops against whitefly attack. The utility of pyriproxyfen in whitefly management was demonstrated based on suppression of embryogenesis and adult formation in *Bemisia tabaci* (Gennadius) (Homoptera: Aleyrodidae).[10] and *Trialeurodes vaporariorum* (Westwood) (Homoptera: Aleyrodidae).[11] In a leaf-disk bioassay using pyriproxyfen on the oblique banded leaf roller, *Choristoneura rosaceana* (Harris) (Lepidoptera: Tortricidae), the LC<sub>50</sub> values for males and females were found to be 2.4 and 4.8 ppm, respectively.[12] Moadeli *et al* (2014)[13] reported that in a leaf dip bioassay with recommended field rate (1000 ppm) of pyriproxyfen, the 1<sup>st</sup> instar larvae of *Spodoptera exigua*, showed delay in larval development and in turn pyriproxyfen prolonged the feeding period and growth. In their study when compared

to control the mean generation time was higher in pyriproxyfen treated insects.

Though insecticides containing pyriproxyfen like 'Knack IGR' are used against a variety of pests, the toxicity of pyriproxyfen is not tested on the larvae of *Spodoptera mauritia*. More over determination of the toxicity of pyriproxyfen will be helpful in determining the sub lethal concentration for use in experiments to study effects on development. In the present study we used 'Knack IGR', a pesticide containing pyriproxyfen, as the active ingredient to test its toxicity on the different larval stages of *Spodoptera mauritia*.

## II. MATERIALS AND METHODS

### A. Collection, Rearing and Maintenance of the Larvae of *Spodoptera Mauritii* Boisid.

The moths were attracted by fluorescent lamps during night. They were collected using an insect sweeping net. The adults were kept in glass chimneys closed at both ends with muslin cloth and fed with 10% solution of honey. The adults lay eggs on the cloth covering the container. The caterpillars were fed with fresh, tender leaves of the grass *Ischaemum aristatum*. Larvae were maintained at room temperature (28°C) with relative humidity of 70-80%

### B. Treatment to Test Toxicity of Pyriproxyfen

From the laboratory culture the 3<sup>rd</sup>, 4<sup>th</sup>, 5<sup>th</sup> and 6<sup>th</sup> instar larvae were sorted out on the basis of moulting marks. Different concentrations of pyriproxyfen in acetone was applied topically on the dorsal side of the 3<sup>rd</sup>, 4<sup>th</sup>, 5<sup>th</sup> & 6<sup>th</sup> instar day 0 larvae of *Spodoptera mauritia* using a Hamilton Micro-Syringe in a total volume of 2µL. An equal volume of acetone was applied in the same manner to the control larvae. At least 3 replicates were done for each experiment and the number of larvae per experiment varied from 10 to 15. Mortality was recorded after 24 hours and from the average percentage mortality for different concentrations of pyriproxyfen, LD<sub>50</sub> value for each instar was calculated from a plot of log concentration versus percentage mortality.

## III. RESULTS

**Table I:** Percentage mortality of 3<sup>rd</sup> 4<sup>th</sup> 5<sup>th</sup> & 6<sup>th</sup> instar larvae of *Spodoptera mauritia* treated with different concentrations of pyriproxyfen

Amount of pyriproxyfen applied/ larva	Average percentage mortality ± SE			
	3 <sup>rd</sup> instar	4 <sup>th</sup> instar	5 <sup>th</sup> instar	6 <sup>th</sup> instar
Control	0	0	0	0
5µg	14.4±2.1	12.5±2.5	.....	.....
10µg	25±5.0	27.5±6.4	.....	.....
25µg	85±2.9	70.8±5.1	15.8±2.0	.....
50µg	96.7±3.3	91.7±4.4	65.8±2.2	.....
100µg	100±0.0	100±0.0	75±2.9	8.13±2.8
125µg	.....	.....	96.7±3.3	13.8±1.3
200µg	.....	.....	.....	32.5±2.5
300µg	.....	.....	.....	45±5.0
400µg	.....	.....	.....	73.3±1.7

### A. Toxicity of Pyriproxyfen to Larvae of *Spodoptera Mauritii*

The average percentage mortality for 3<sup>rd</sup>, 4<sup>th</sup>, 5<sup>th</sup> & 6<sup>th</sup> instar larvae of *Spodoptera mauritia* treated with different concentrations of pyriproxyfen was calculated (Table 1.) With increase in concentration of pyriproxyfen, the mortality increased in all the instars of larvae tested.

### B. Calculation of LD<sub>50</sub> value

The LD<sub>50</sub> value (24 hours) of pyriproxyfen for the 3<sup>rd</sup>, 4<sup>th</sup>, 5<sup>th</sup> & 6<sup>th</sup> instar larvae of *Spodoptera mauritia* was found to be 14.13±2.67, 15.85±3.67, 39.81±2.61 and 316.20±2.64 µg/larvae respectively (Table 2)

**Table 2:** LD<sub>50</sub> value (24 hours) of pyriproxyfen for 3<sup>rd</sup> 4<sup>th</sup> 5<sup>th</sup> & 6<sup>th</sup> instar larvae of *Spodoptera mauritia*

SL. NO.	LARVAL NSTAR	LD <sub>50</sub> VALUE (µg) (MEAN ±SE)
1	THIRD	14.13±2.67
2	FOURTH	15.85±3.67
3	FIFTH	39.81±2.61
4	SIXTH	316.20±2.64

## IV. DISCUSSION

When the 3<sup>rd</sup> instar day 0 larvae were treated with 5, 10, 25, 50 and 100 µg per larva of pyriproxyfen, the average percentage mortality after 24 hours was found to be 14.38±2.13, 25±5.0, 85±2.88, 96.67±3.33 and 100±0.0 percentage respectively. The LD<sub>50</sub> 24 hours, calculated for the 3<sup>rd</sup> instar larvae is 14.13±2.67µg (Table: 2). In the case of *Plutella xylostella* (L.) (Lepidoptera: Plutellidae) 3<sup>rd</sup> instar larvae, the reported LC<sub>50</sub> value based on a leaf dip bioassay is 1.223 g L<sup>-1</sup>. [14]. When pyriproxifen was incorporated in the artificial diet of ten-day-old larvae of Indian meal moth *Plodia interpunctella* (Lepidoptera: Pyralidae) at 0.02, 0.04, 0.08, 0.16, and 0.3 ppm concentrations, it resulted in increased duration of the larval period till the emergence of adults, decreased adult longevity and significant reduction in mean number of eggs laid by adults compared to the controls[15]

In the case of 4<sup>th</sup> instar day 0 larvae treated with pyriproxyfen, 5, 10, 25, 50 and 100µg per larva, the average percentage mortality after 24 hour was found to be 12.5±2.50, 27.5±6.37, 70.83±5.07, 91.67±4.41 and 100±0.0 percentages respectively. The LD<sub>50</sub> 24 hour for the 4<sup>th</sup> instar larvae is 15.85±3.67µg/larva. In the freshly moulted fourth instar larvae of citrus swallowtail *Papilio demoleus* (Lepidoptera: Papilionidae) topical administration of pyriproxyfen (7.5, 15, 30 and 60 µg/1µl/larva) induced a delay in larval– larval ecdysis and subsequent larval–pupal ecdysis. It is also reported that the treatment reduced the frequency of pupation, increased mortality and ecdysial failure, and inhibited adult emergence. [16]

For the 5<sup>th</sup> instar day 0 larvae there was no mortality for 5µg and 10µg pyriproxyfen per larva. Thus higher amount of pyriproxyfen was applied. The average percentage mortality after 24 hours for 25, 50, 100 and 125µg of pyriproxyfen per larva were 15.83±2.01, 65.83±2.21, 75±2.89 and 96.67±3.33 percentage respectively. The



LD<sub>50</sub> calculated based on this percentage mortality was found to be 39.81±2.61µg/larva.

Sixth instar day 0 larvae when treated with 100, 200, 300 & 400µg of pyriproxyfen per larvae, the mortality was found to be 8.13±2.77, 13.75±1.25, 32.5±2.5, 45±5.0 and 73±1.67 percentage respectively. On calculation the LD<sub>50</sub> value was found to be 316.20±2.64µg/larvae. Singh et al (2015)[17] topically administrated sub-lethal doses (0.5, 1.0, 2.5 & 5µg/ µl/larvae) of pyriproxyfen and difenolan on the 6<sup>th</sup> instar larvae of *Spodoptera litura* (Lepidoptera: Noctuidae) and reported that these JHAs severely hampered the metamorphosis and development with prolonged larval duration, mortality, ecdysial failure, formation of larval - pupal mosaics, reduced pupation and formation of abnormal pupae, complete suppression of adult emergence and production of adultoids. Significant differences in number and hatchability of eggs, wing abnormalities and morphological ovarian abnormalities were observed when pyriproxyfen was topically applied to *Spodoptera litura* (Lepidoptera: Noctuidae). Application of 0.1 ng of pyriproxyfen to day-1 female pupae and 0.125 µg to day-0 6th stadium larvae reduced the total number of eggs oviposited and hatchability of eggs. Day-1 pupal stage treated with 0.3 ng of pyriproxyfen showed wing abnormalities and about 40% of female adults showed morphological ovarian abnormalities. [18]

As pyriproxyfen is effective at lower concentrations to disrupt the larval development leading to failure in healthy adult emergence, most of the studies are concentrated on sublethal effects. In this study we showed that pyriproxyfen at relatively higher concentration causes death of the larvae in 24 hours after application. The results can be extrapolated to find sub-lethal concentration such as LD<sub>10</sub> or other sub lethal concentrations for *S. mauritia* for experiments to study its effect on development. Studies are ongoing to find effect of sub lethal concentration of pyriproxifen on development of *Spodoptera mauritia*.

## V. CONCLUSIONS

At higher concentrations, pyriproxyfen caused the death of the larvae of *S.mauritia* after 24 hours and with increase in concentration of pyriproxyfen, the mortality also increased in 3<sup>rd</sup> 4<sup>th</sup> 5<sup>th</sup> and 6<sup>th</sup> instar larvae. The LD<sub>50</sub> value of pyriproxyfen for the 3<sup>rd</sup>, 4<sup>th</sup>, 5<sup>th</sup> and 6<sup>th</sup> instar larvae of *Spodoptera mauritia* was found to be 14.13±2.67, 15.85±3.67, 39.81±2.61 and 316.20±2.64 µg/larvae respectively.

## VI. ACKNOWLEDGMENTS

University Grants Commission, New Delhi for financial support and UGC Special Assistance Programme (SAP), Department of Zoology for the facilities.

## REFERENCES

- [1] B. V. David, "The swarming caterpillar or armyworm, *Spodoptera mauritia*" *Journal of Insect Physiology* 37(2), 2004, pp 87-93

- [2] A. Retnakaran, J. Granett and T. Ennis "Insect Growth Regulators" In *Comprehensive Insect Physiology, Biochemistry and Pharmacology* (Ed. by Kerkut, G.A. and Gilbert, L.I.). Pergamon Press, Oxford. 1985, 12 pp.529-601
- [3] B. Darvas, L. Varjas, "Insect Growth Regulators In armored scale insect, their biology natural enemies control" (Ed. by Rosen, D). Elsevier science publishers, B.V. Amsterdam, Vol.B, 1990, pp.393-408.
- [4] V.S.K. Nair, "Applied potential of insect hormone research perspectives and prospects" In *Chemical Ecology of Phytophagous Insects* (Ed. Ananthakrishnan, T.N. and Raman, A). Oxford-IBH Publ. Co., New Delhi, 1993, pp89-104.
- [5] B.A. Peleg, "Effect of a new insect growth regulator", Rol3-5223 on scale insects. *Phytoparasitology*, vol.10, 1982, pp.27-31.
- [6] C.A. Henrick, G.B. Staal, J.B. Siddal, "Alkyl3,7,11-trimethyl-2,4-dodecadienoates, a new class of potent insect growth regulators with juvenile hormone activity" *Journal of Agriculture and Food Chemistry*, vol.21, 1973, pp.354-359.
- [7] J.L. Krysan, "Fenoxycarb and diapause: a possible method of control for pear psylla (Homoptera: Psyllidae)", *Journal of Economic Entomology*, vol.83, 1990, pp.293-299.
- [8] J. Miyamoto, M. Hirano, Y. Takimoto, M. Hatakoshi, "Insect growth regulator for pest control, with emphasis on juvenile hormone analogs—present status and future prospects" In: Duke, S.O., Menn, J.J., Plimmer, J.R. (Eds.), *Pest Control with Enhanced Environmental Safety*, vol.524, American Chemical Society Symposium, Washington, DC, 1993, pp.144-168.
- [9] P.C. Ellsworth, J.W. Diehl, I.W. Kirk, T.J. Henneberry, "Bemisia growth regulators: large-scale evaluation" In: Dugger, P., Richter, D. (Eds.), *Proceedings of the Beltwide Cotton Conferences. Cotton Insect Research and Control Conference*, Nashville, TN, 1997, pp.922-929.
- [10] I. Ishaaya, A.R. Horowitz, "Novel phenoxy juvenile hormone analog (pyriproxyfen) suppresses embryogenesis and adult emergence of sweetpotato whitefly (Homoptera: Aleyrodidae)", *J. Econ. Entomol.*, vol.85, 1992, pp.2113-2117.
- [11] I. Ishaaya, A. De Cock, D. Degheele, "Pyriproxyfen, a potent suppressor of egg hatch and adult formation of the greenhouse whitefly (Homoptera: Aleyrodidae)", *J. Econ. Entomol.*, vol.87, 1994, pp.1185-1189.
- [12] A.A. Sial, J.F. Brunner, "Lethal and sublethal effects of an insect growth regulator, pyriproxyfen, on obliquebanded leafroller (Lepidoptera: Tortricidae)" vol. 103(2), *J Econ Entomol*, 2010, pp. 340-7.
- [13] T. Moadeli, M.J. Hejazi, Gh. Golmohammadi, "Lethal Effects of Pyriproxyfen, Spinosad, and Indoxacarb and Sublethal Effects of Pyriproxyfen on the 1st Instars Larvae of Beet Armyworm, *Spodoptera exigua* Hübner (Lepidoptera: Noctuidae) in the Laboratory", *J. Agr. Sci. Tech.*, vol. 16, 2014, pp.1217-1227.
- [14] M. Mahmoudvand, S. Moharrampour, and M. Iranshahi, "Effects of Pyriproxyfen on Life Table Indices of *Plutella xylostella* in Multigenerations", *Hindawi Publishing Corporation, Psyche*, Volume 2015 (2015), 2015, 7 pages.
- [15] A. Ghasemi, J. J. Sendi and M. Ghadamyari, "Physiological and biochemical effect of pyriproxyfen on indian meal moth *Plodia interpunctella* (hübner) (Lepidoptera: Pyralidae)", *Journal of plant protection research* vol. 50, no. 4 (2010).
- [16] S. Singh and K. Kumar, "Effect of the juvenile hormone agonist pyriproxyfen on larval and pupal development of the citrus swallowtail *Papilio demoleus* (Lepidoptera: Papilionidae)" *International Journal of Tropical Insect Science*, vol. 31(3), 2011, pp. 192-198.
- [17] S. Singh and K. Kumar, "Comparative efficacy of phenoxy derivative JHAs Pyriproxyfen and Difenoalan against polyphagous pest *Spodoptera litura* (Fabricius) (Noctuidae: Lepidoptera)", *Phytoparasitica*, vol. 43, 2015, pp. 553-563.
- [18] M. Nomura and T. Miyata, "Effects of Pyriproxyfen, Insect Growth Regulator on Reproduction of Common Cutworm, *Spodoptera litura* (Fabricius) (Lepidoptera: Noctuidae)", *Japanese Journal of Applied Entomology and Zoology*, vol. 44(2), pp 81-88.

**AUTHORS' PROFILES**

**Dr. Kannan Vadakkadath Meethal** hails from Ulleary village in Kozikode District of Kerala, India. He completed M.Sc Zoology from University of Calicut and Ph.D from Sree Chitra Tirunal Institute for Medical Sciences and Technology, Trivandrum. He completed his Post Doctoral Training from University of Michigan,



**Mrs. Resmitha C.** hails from Thazhecode village of Kozhikode District, Kerala, India and completed M.Sc in Zoology from University of Calicut. She is currently pursuing Ph.D in Zoology and working as a research fellow with Dr. Kannan Vadakkadath Meethal in University of Calicut.